



Salmon Stream Temperature Monitoring In the Chilkat & Chilkoot Watersheds

Summary Report – December 2022

Prepared by the Takshanuk Watershed Council

Stacie Evans, Science Director stacie@takshanuk.org

Derek Poinsette, Executive Director derek@takshanuk.org



This report provides an update in progress and summary statistics for water temperature sites monitored by Takshanuk Watershed Council. Monitoring at these sites is part of an ongoing collaboration with the Southeast Alaska Freshwater Monitoring Network. Funding was provided by Chilkat Indian Village, National Fish and Wildlife Foundation, US Fish and Wildlife Service, and Patagonia.

Contents

Background	1
Methods.....	2
Site Selection.....	2
TWC data logger locations.	3
Data Collection.....	5
Data Management and Summarization	6
Results.....	7
West Lake Creek.....	8
Herman Creek	9
Clear Creek	10
Little Salmon River	11
14-Mile Creek.....	12
Lower Kelsall River	13
Lower Chilkoot River	14
Mink Creek	15
Tahini River	16
Jones Point (Chilkat River)	16
Sarah Creek	17
JKHC (Chilkat River)	17
Chilkat Lake Weir	18
Upper Chilkoot River	18
Northeast Creek	19
Davis Creek.....	19
Assignation Creek.....	20
Mule Meadows	20
Upper Chilkat Slough.....	21
MWMTs and MWATs.....	22
Exceedances.....	23
Yearly Comparison	24
Discussion.....	25
Literature Cited	27

Background

The Chilkat and Chilkoot watersheds are two of the most productive salmon systems on the west coast of North America. They host significant wild runs of all five species of Pacific salmon, as well as steelhead, Dolly Varden, coastal cutthroat trout, and eulachon. This abundance has supported the Jilḱáat Ƴwáan and Jilḱoot Ƴwáan since time immemorial, and continues to support them today. In addition, the commercial fishing industry accounts for 15% of local incomes and virtually all residents benefit from subsistence and traditional hunting, fishing, and gathering (adfg.alaska.gov). Beyond economic benefits, the practice of subsistence food collection and preservation promotes family ties, community resiliency, and passes on tradition and cultural identity to future generations. Although local freshwater habitats are relatively intact, they are nonetheless facing a significant near-term threat: the unpredictable effects of climate change.

The climate is changing rapidly in the North. Over the next 50 years, Southeast Alaska can expect to see an increase in mean annual air temperature of 2 to 4 °C, as well as drier summers and wetter autumns (Schoen et al. 2017). These factors, along with watershed characteristics and stream morphology, drive the potential for widespread and rapid changes in stream temperature and the local aquatic environment (Kovach et al. 2015, Shanley et al. 2015). The region is also heavily glaciated, and we expect to see significant changes in both glacial runoff and precipitation patterns, which will greatly impact habitat. The generalized effects of water temperature on the health, growth, and behavior of aquatic life, especially salmon, are well studied and well understood (Steel et al. 2012, Taylor 2008, Webb et. al 2008). What are not well documented are the specific local processes that are occurring on the landscape and within the habitat, especially over a time scale of years and decades.

The immediate purpose of this project is to develop a baseline, year-round temperature dataset to document the current condition of sites within the Chilkoot and Chilkat watersheds. Over time we aim to generate a long term dataset to discern thermal trends across both watersheds. In addition, these data contribute to collaborative efforts supported by The Southeast Alaska Freshwater Temperature Monitoring Network to standardize, share, and interpret data across the region, and may also be used to create predictive models to inform management decisions regarding aquatic habitat (Bellmore and Winfree 2019).

This report is intended to provide an update in progress for this project and to help identify areas to expand our monitoring efforts. The data summaries are also useful for identifying sites that exceed ADEC maximum temperature limits for aquatic life and may be experiencing, or at risk of experiencing, habitat degradation for salmon.

Methods

Methods for data collection, management, and summary were adapted from *Stream Temperature Data Collection Standards and Protocol for Alaska* (Mauger et al. 2014). Refer to that manual for a more detailed description of methods, protocols, and equipment.

Site Selection

Sites were selected to cover a broad range of hydrological and physical characteristics, and all sites contained salmon habitat (Table 2). Relative ease of access was an important factor in determining where to deploy data loggers, although off-road and remote sites were also selected. Instantaneous temperature measurements were taken across the stream to verify that sites were well mixed before deploying loggers, and loggers were placed in locations where they were unlikely to become dewatered due to low stream flow.

Table 1. Takshanuk Watershed Council data logger locations and activity periods.

Site	Latitude	Longitude	Date Deployed	Latest Download Date*
West Lake Creek	59.33748	-135.59080	5/28/2016	9/7/2022
Herman Creek	59.41326	-136.06848	5/29/2016	9/8/2022
Clear Creek	59.38226	-136.06812	5/29/2016	9/8/2022
Little Salmon River	59.38322	-136.07504	9/29/2017	9/8/2022
14-Mile Creek	59.32871	-135.74188	5/8/2018	11/17/2022
Lower Kelsall River	59.54015	-136.10397	9/18/2018	9/21/2022
Lower Chilkoot River	59.33070	-135.55608	11/21/2018	10/14/2022
Mink Creek	59.15719	-135.35882	7/17/2019	9/15/2022
Tahini River	59.64146	-135.99401	9/11/2019	9/12/2022
Jones Point	59.22998	-135.48507	4/15/2020	9/15/2022
Sarah Creek	59.42850	-136.32688	6/5/2020	10/7/2022
JKHC	59.39814	-135.88493	6/3/2021	11/17/2022
Chilkat Lake Weir	59.36206	-135.93813	6/8/2021	10/25/2022
Upper Chilkoot River	59.37222	-135.63531	8/4/2021	10/14/2022
Northeast Creek	59.37538	-135.61290	8/4/2021	10/14/2022
Davis Creek	59.36000	-135.61791	8/4/2021	10/14/2022
Assignment Creek	59.56431	-135.99688	8/16/2021	9/12/2022
Mule Meadows	59.54371	-136.05695	8/16/2021	9/12/2022
Upper Chilkat Slough	59.29452	-135.82191	6/9/2022	10/25/2022

*Data loggers are still active at each site. The "Latest Download Date" is the most recent date that data were downloaded and summarized for this report.

Takshanuk Watershed Council
Temperature Monitoring Sites
2021

Map showing the Takshanuk Watershed Council Temperature Monitoring Sites for 2021. The map displays the watershed area, including major rivers (Chilkat River, Chilkoot River, Tahini River, Flenier River, Takhin River, Skagway River, Klondike River, Jewa River), creeks (Sarah Creek, Herman Creek, Little Salmon, Clear Creek, Northeast Creek, Davis Creek, West Lake Creek, Lower Chilkoot River, Narrows Creek, Upper Chilkat Slough, Mink Creek), and lakes (Chilkat Lake, Upper Chilkoot Lake). The map also shows the border between Canada and the United States, and the location of Klukwan and Haines. A north arrow and a scale bar (0 to 30 miles) are included.

Monitoring sites marked on the map include:

- Tahini River
- Assignment Creek
- Mule Meadows
- Lower Kelsall River
- Sarah Creek
- Herman Creek
- Little Salmon
- Clear Creek
- Chilkat Lake Weir
- Upper Chilkoot River
- Northeast Creek
- Davis Creek
- West Lake Creek
- Lower Chilkoot River
- Narrows Creek
- Upper Chilkat Slough
- Jones Point
- Mink Creek

Geographic features and locations shown include:

- Canada / United States border
- Klukwan
- Haines
- Chilkat Inlet
- Chilkat State Park
- Takshanuk Mountains
- Takhinsha Mountains
- Henry Clay
- Skagway
- Klondike Hwy
- Chilkat River
- Chilkoot River
- Tahini River
- Flenier River
- Takhin River
- Skagway River
- Klondike River
- Jewa River
- Chilkat Lake
- Upper Chilkoot Lake
- Chilkat Lake Weir
- Upper Chilkoot River
- Northeast Creek
- Davis Creek
- West Lake Creek
- Lower Chilkoot River
- Narrows Creek
- Upper Chilkat Slough
- Jones Point
- Mink Creek
- Chilkat Inlet
- Chilkat State Park
- Takshanuk Mountains
- Takhinsha Mountains
- Henry Clay
- Skagway
- Klondike Hwy
- Chilkat River
- Chilkoot River
- Tahini River
- Flenier River
- Takhin River
- Skagway River
- Klondike River
- Jewa River
- Chilkat Lake
- Upper Chilkoot Lake
- Chilkat Lake Weir
- Upper Chilkoot River
- Northeast Creek
- Davis Creek
- West Lake Creek
- Lower Chilkoot River
- Narrows Creek
- Upper Chilkat Slough
- Jones Point
- Mink Creek
- Chilkat Inlet
- Chilkat State Park



Stacie Evans (TWC Science Director) and Richard Chapell (TWC Board Member) headed back from the Tahini site. Photo by Derek Poinsette.



Apparently there's more than one way to test the water. Chilkoot Lake. Photo by Derek Poinsette.



Preparing to swap data loggers at the West Lake Creek Site. Photo by Derek Poinsette.

Data Collection

HOBO Water Temp Pro v2 data loggers were checked for $\pm 0.25^{\circ}\text{C}$ accuracy against a National Institute of Science and Technology (NIST) certified thermometer before each deployment in the field. Data loggers were programmed to record temperature every 30 minutes, except for at Herman Creek, Clear Creek, and West Lake Creek where loggers were programmed to record temperature every one hour from May 2016 to November 2018.

Three data loggers were deployed at each site – two in-stream to record water temperature, and one out of stream to record air temperature. Each logger was secured with zip ties in a protective PVC housing. In-stream loggers were attached to heavy chain (to weigh the loggers down) and secured with a loop of cable and clamps to a large rock or sturdy object on the stream bank. Air temperature loggers were hung in a shady location in nearby vegetation.

Data loggers were retrieved and fresh loggers were deployed every spring and fall to the extent possible. Loggers were checked again for accuracy and battery power upon retrieval. Sites were checked opportunistically to ensure that loggers were in place, operating, and free from sedimentation.



Derek Poinsette (TWC Executive Director) and Richard Chapell (ADF&G) deploying data loggers at the Tahini Site.
Photo by Jenn Hamblen

Data Management and Summarization

Data were downloaded from each data logger using HOBOWare software. Hobo graph outputs were examined for erroneous data due to dewatering or sedimentation before data were exported to comma-separated values (CSV) files for backup and Excel files for summarization. Pre deployment, post retrieval, and erroneous data were removed from Excel spreadsheets. The remaining data (for days containing at least 90% of the daily measurements) were organized in pivot tables and summarized into daily minimum, maximum, and mean water temperatures. Maximum 7-day rolling average (MWAT) and maximum 7-day rolling maximum (MWMT) temperatures were also summarized for summer months at each site. MWMT was calculated by selecting the yearly maximum from weekly averages of maximum daily temperatures and MWAT was calculated by selecting the yearly maximum from weekly averages of average daily temperatures.

Daily temperatures, MWMTs, and MWATs at each site were compared to the ADEC water temperature criteria for aquatic life in freshwater (ADEC 2020). These standards are as follows:

May not exceed 20°C at any time. The following maximum temperatures may not be exceeded, where applicable:

Migration routes	15°C
Spawning areas	13°C
Rearing areas	15°C
Egg and fry incubation	13°C

Additional years of data are needed to consider inter-annual variability at some sites, however, daily mean temperatures were compared by year for sites with at least three years of useable data. Average July and August temperatures were also compared by year for those sites.

Results


Data loggers were deployed at 19 sites, beginning with three sites established in May 2016 (Table 1). Sites were subsequently added, with the latest site established in June 2022 (Table 1). Gaps in the data represent periods when data loggers were lost, or when erroneous data caused by dewatering or sedimentation were removed. See figure captions for details at each site.

Summary graphs with daily minimum, maximum, and mean temperatures are presented below for each site. Sites that exceeded ADEC water temperature criteria for aquatic life in freshwater are reported in Table 2. Temperatures above 20°C were only recorded at the Chilkat Weir site. The number of days in which temperatures exceeded ADEC standards at sites is reported in Table 3 by year. Summary graphs are also included for yearly maximum 7-day rolling maximum temperatures (MWMTs) and maximum 7-day rolling average temperatures (MWATs). Exceedances with these metrics are included in Table 2. MWMTs and MWATs are useful metrics because they describe thermal conditions that are experienced by fish over an extended period of time (a week), rather than discrete measurements which may only represent short spikes in temperature.

Yearly comparisons of daily mean temperatures are included for sites where at least three complete years of data were available (West Lake Creek, Herman Creek, Clear Creek, Little Salmon River, 14-Mile Creek, Lower Kelsall River, Lower Chilkoot River, and Mink Creek). Mean July/August temperatures at these sites are presented in Figure 30 by year.


West Lake Creek

Figure 1. Daily minimum, maximum, and mean temperatures at the West Lake Creek site from 5/28/2016 to 9/7/2022.



*13°C = ADEC maximum temperature limit for spawning areas and egg/fry incubation. 15°C = ADEC maximum temperature limit for migration routes and rearing areas.

Figure 2. Yearly comparison of daily mean temperatures at the West Lake Creek site.



Herman Creek

Figure 3. Daily minimum, maximum, and mean temperatures at the Herman Creek site from 5/28/2016 to 9/8/2022. Data are not included from 12/3/2021 to 5/8/2021; a period when data loggers were affected by sediment.





Figure 4. Yearly comparison of daily mean temperatures at the Herman Creek site.



Clear Creek

Figure 5. Daily minimum, maximum, and mean temperatures at the Clear Creek site from 5/29/2016 to 9/8/2022.






Figure 6. Yearly comparison of daily mean temperatures at the Clear Creek site.




Little Salmon River

Figure 7. Daily minimum, maximum, and mean temperatures at the Little Salmon River site from 9/29/2017 to 9/8/2022. Data are not included from 5/27/2020 to 6/2/2021; a period when one logger washed away and the other logger malfunctioned.



*13°C=ADEC maximum temperature limit for spawning areas and egg/fry incubation. 15°C=ADEC maximum temperature limit for migration and rearing areas.

Figure 8. Yearly comparison of daily mean temperatures at the Little Salmon River site.



14-Mile Creek

Figure 9. Daily minimum, maximum, and mean temperatures at the 14-Mile Creek site from 5/8/2018 to 11/17/2022.






Figure 10. Yearly comparison of daily mean temperatures at the 14-Mile Creek site.




Lower Kelsall River

Figure 11. Daily minimum, maximum, and mean temperatures at the Lower Kelsall River site from 9/18/2018 to 9/21/2022. Data are not included from 10/3/2019 to 5/22/2020 or 9/13/2020 to 6/8/2021; periods when loggers washed away or were dewatered.




*13°C=ADEC maximum temperature limit for spawning areas and egg/fry incubation. 15°C=ADEC maximum temperature limit for migration and rearing areas.

Figure 12. Yearly comparison of daily mean temperatures at the Lower Kelsall River site.




Lower Chilkoot River

Figure 13. Daily minimum, maximum, and mean temperatures at the Lower Chilkoot River site from 11/21/2018 to 10/14/2022. Data are not included from 10/28/2020 to 1/19/2021; a period when the cable rusted through and loggers were removed until replacement was possible.




*13°C=ADEC maximum temperature limit for spawning areas and egg/fry incubation. 15°C=ADEC maximum temperature limit for migration and rearing areas.

Figure 14. Yearly comparison of daily mean temperatures at the Lower Chilkoot River site.




Mink Creek

Figure 15. Daily minimum, maximum, and mean temperatures at the Mink Creek site from 7/17/2019 to 9/15/2022.




*13°C=ADEC maximum temperature limit for spawning areas and egg/fry incubation. 15°C=ADEC maximum temperature limit for migration and rearing areas.

Figure 16. Yearly comparison of daily mean temperatures at the Lower Kelsall River site.



Tahini River


Figure 17. Daily minimum, maximum, and mean temperatures at the Tahini River site from 9/11/2019 to 9/12/2022. Data are not included from 5/22/2020 to 6/8/2021; a period when both data loggers washed away.



*13°C=ADEC maximum temperature limit for spawning areas and egg/fry incubation. 15°C=ADEC maximum temperature limit for migration and rearing areas.

Jones Point (Chilkat River)


Figure 18. Daily minimum, maximum, and mean temperatures at the Jones Point site from 4/15/2020 to 9/15/2022. Data are not available from 9/22/2020 to 3/24/2022 because loggers were buried and irretrievable. Loggers were moved to a better site just downstream.



*13°C=ADEC maximum temperature limit for spawning areas and egg/fry incubation. 15°C=ADEC maximum temperature limit for migration and rearing areas.


Sarah Creek

Figure 19. Daily minimum, maximum, and mean temperatures at the Sarah Creek site from 6/6/2020 to 10/7/2022. Data from 10/21/2020 to 10/17/2021 are not available because loggers washed away.



JKHC (Chilkat River)


Figure 20. Daily minimum, maximum, and mean temperatures at the JKHC site from 6/3/2021 to 11/17/2022.



*13°C=ADEC maximum temperature limit for spawning areas and egg/fry incubation. 15°C=ADEC maximum temperature limit for migration and rearing areas.

Chilkat Lake Weir


Figure 21. Daily minimum, maximum, and mean temperatures at the Chilkat Lake Weir site from 6/8/2021 to 10/25/2022.



*13°C=ADEC maximum temperature limit for spawning areas and egg/fry incubation. 15°C=ADEC maximum temperature limit for migration and rearing areas. 20°C=ADEC maximum temperature which may not be exceeded at any time.

Upper Chilkoot River


Figure 22. Daily minimum, maximum, and mean temperatures at the Upper Chilkoot River site from 5/12/2022 to 10/14/2022.



*13°C=ADEC maximum temperature limit for spawning areas and egg/fry incubation. 15°C=ADEC maximum temperature limit for migration and rearing areas.

Northeast Creek


Figure 23. Daily minimum, maximum, and mean temperatures at the Northeast Creek site from 8/5/2021 to 10/14/2022.



*13°C=ADEC maximum temperature limit for spawning areas and egg/fry incubation. 15°C=ADEC maximum temperature limit for migration and rearing areas.


Davis Creek

Figure 24. Daily minimum, maximum, and mean temperatures at the Davis Creek site from 8/4/2021 to 10/14/2022. Data are not available from 11/8/2021 to 5/11/2022 due to dewatering.



Assignation Creek


Figure 25. Daily minimum, maximum, and mean temperatures at the Assignation Creek site from 8/17/2022 to 9/3/2022.



*13°C=ADEC maximum temperature limit for spawning areas and egg/fry incubation. 15°C=ADEC maximum temperature limit for migration and rearing areas.

Mule Meadows


Figure 26. Daily minimum, maximum, and mean temperatures at the Mule Meadows site from 8/17/2022 to 9/3/2022.



*13°C=ADEC maximum temperature limit for spawning areas and egg/fry incubation. 15°C=ADEC maximum temperature limit for migration and rearing areas.

Upper Chilkat Slough


Figure 27. Daily minimum, maximum, and mean temperatures at the Upper Chilkat Slough site from 6/9/2022 to 10/25/2022.



*13°C=ADEC maximum temperature limit for spawning areas and egg/fry incubation. 15°C=ADEC maximum temperature limit for migration and rearing areas.


MWMTs and MWATs

Figure 28. Summer maximum weekly maximum temperature (MWMT) by site. Data from June 1 to September 30 are included.



*13°C=ADEC maximum temperature limit for spawning areas and egg/fry incubation. 15°C=ADEC maximum temperature limit for migration and rearing areas. 20°C=ADEC maximum temperature which may not be exceeded at any time.

Figure 29. Summer maximum weekly average temperature (MWAT) by site.



*13°C=ADEC maximum temperature limit for spawning areas and egg/fry incubation. 15°C=ADEC maximum temperature limit for migration and rearing areas. 20°C=ADEC maximum temperature which may not be exceeded at any time.

Exceedances

Table 2. Maximum temperature recorded at each site. Sites where temperatures exceeded ADEC water temperature limits for aquatic life in freshwater are noted.

Site	Max Temp°C	Date Max Recorded	Salmon Species Present ¹	Max Temp Exceeded			MWMT Exceeded			MWAT Exceeded	
				13°C	15°C	20°C	13°C	15°C	20°C	13°C	15°C
West Lake Creek	12.654	8/7/2019	S								
Herman Creek	7.845	8/17/2022	CH, CO, P, S								
Clear Creek	7.167	7/4/2019	CO, S								
Little Salmon River	13.810	8/6/2019	CH, CO, S	X	X		X				
14-Mile Creek	14.361	7/21/2018	CH, CO, P	X			X				
Lower Kelsall River	17.772	7/5/2019	CH, CO, K, P, S	X	X		X	X		X	X
Lower Chilkoot River	18.438	7/6/2019	CH, CO, P, S	X	X		X	X		X	
Mink Creek	15.963	8/7/2019	CH, CO	X	X		X				
Tahini River	14.098	8/1/2021	CO, K, S	X							
Jones Point	17.391	5/19/2022	CH, CO, K, P, S	X	X		X	X			
Sarah Creek	10.345	7/30/2020	CO								
JKHC	11.516	7/3/2022	CH, CO, K, P, S								
Chilkat Weir	22.345	6/28/2022	CH, CO, S	X	X	X	X	X	X	X	X
Upper Chilkoot River	11.540	7/4/2022	CH, CO, P, S								
Northeast Creek	11.904	7/3/2022	CO								
Davis Creek	9.485	8/21/2022	S								
Assignment Creek	12.630	8/4/2022	CO, K, P								
Mule Meadows	10.492	9/19/2021	CO, S								
Upper Chilkat Lake	9.706	8/25/2022	CO, S								

¹According to the Alaska Department of Fish and Game Anadromous Waters Catalogue (ADF&G 2021). CH=Chum, CO=Coho, K=Chinook, P=Pink, S=Sockeye.

Eight out of nineteen sites exceeded the lower limit of 13°C for spawning areas and egg/fry incubation. The MWMT exceeded 13°C at seven of these sites. Six sites exceeded the middle limit of 15°C for migration routes and rearing areas, and the MWMT exceeded this limit at four of these sites. The MWAT exceeded 13°C at Lower Chilkoot, Lower Kelsall, and Chilkat Weir, and it also exceeded 15°C at Lower Kelsall, and Chilkat Weir. The maximum temperature and the MWMT at Chilkat Weir exceeded 20°C, which may not be exceeded at any time (ADEC 2020).

Maximum stream temperatures varied broadly across sites with a low maximum of 7.167°C at Clear Creek and a high maximum of 22.345°C at Chilkat Weir. Herman Creek and Clear Creek are largely fed by groundwater, which likely contributes to less temperature fluctuation compared to other sites.

Table 3. Number of days per year in which water temperatures exceeded ADEC limits for aquatic life in freshwater. Dashes indicate that data were not collected for that year. Data loggers were not deployed before 2018 at sites where exceedances occurred.


Site	2018		2019		2020		2021			2022		
	#Days Exceeded 13°C*	# Days Exceeded 15°C*	#Days Exceeded 13°C*	# Days Exceeded 15°C*	#Days Exceeded 13°C*	# Days Exceeded 15°C*	#Days Exceeded 13°C*	# Days Exceeded 15°C*	# Days Exceeded 20°C*	#Days Exceeded 13°C*	# Days Exceeded 15°C*	# Days Exceeded 20°C*
Little Salmon	6	0	13	0	-	-	11	2	0	0	0	0
14-Mile Creek	12	0	8	0	2	0	3	0	0	0	0	0
Lower Kelsall	-	-	49	28	7	2	17	2	0	28	5	0
Lower Chilkoot	-	-	41	6	0	0	1	0	0	1	0	0
Mink Creek	-	-	30+ [†]	3+ [†]	17	0	11	0	0	38	5	0
Tahini	-	-	-	-	-	-	2	0	0	1	0	0
Jones Point	-	-	-	-	60	22	-	-	-	65	30	0
Chilkat Weir	-	-	-	-	-	-	82	68	2	108	79	10

*13°C=ADEC maximum temperature limit for spawning areas and egg/fry incubation. 15°C=ADEC maximum temperature limit for migration and rearing areas.

[†]Data loggers were deployed July 17, 2019. It is likely that additional exceedances occurred at Mink Creek prior to that date in 2019.

Yearly Comparison

Figure 30. Average stream temperature from 1-July to 31-August by site. Only sites with at least three years of data for July and August are included (West Lake Creek, Herman Creek, Clear Creek, Little Salmon, 14-Mile Creek, Lower Kelsall River, Lower Chilkoot River, and Mink Creek).



Discussion

This stream temperature monitoring program has initiated a valuable thermal dataset for the Chilkat and Chilkoot watersheds. While no formal analyses were performed for this report, summary statistics indicate that much of the salmon habitat associated with these sites is vulnerable to rising temperatures due to climate change. More descriptive, spatial, and temporal data are necessary to better quantify current effects and predict how changes may occur under likely climate scenarios.

Exceedances of ADEC stream temperature limits under current conditions suggest that some salmon habitat is already experiencing degradation due to warm temperatures in the Chilkat and Chilkoot watersheds (Table 3). This is perhaps most concerning at the Chilkat Weir, Lower Chilkoot River, Lower Kelsall River, and Jones Point (Chilkat River); sites where maximum temperatures were the highest, exceedances the most frequent, and most or all species of Pacific salmon are present (Table 2 and 3). On the other hand, these watersheds appear to be faring better than other regions, such as southcentral Alaska, where data loggers in the Deshka River recorded a high of 27.6°C on July 7, 2019. Systems with abundant groundwater inputs, like the Chilkat watershed, could provide important refugia for Pacific salmon as temperatures continue to rise.

Air temperature data are available for each site, but have not yet been summarized for this project. These data will be used in future analyses to determine the sensitivity of stream temperature to local air temperature. This metric is necessary to predict how streams may respond to likely climate scenarios and to determine which sites are at the highest risk. For example, sites that are already warm and also have a high sensitivity to air temperature will be the most vulnerable to impacts due to climate change. Conversely, colder streams with low sensitivity to air temperature are more likely to provide thermal refugia for salmon over time (Mauger 2013).

In addition, characteristic data for each site can be used to determine variables that influence thermal response and to what degree. Predictor variables help managers focus their response to climate change on specific actions with a high degree of resolution within the watershed. Manageable, small-scale strategies, like restoring riparian vegetation to shade a stream, or improving fish passage to less vulnerable habitat, will contribute to the overall resiliency of salmon populations (Mauger et al. 2017). A targeted approach will similarly allow managers to prioritize limited financial resources and increase efficiency of conservation actions (Isaak and Rieman 2013).

While further analyses will be helpful, the preliminary assessment of this dataset can be used to identify current areas of concern. For example, Chilkat chinook salmon were designated a stock of conservation concern by the Alaska Department of Fish and Game in 2017 due to frequent low returns since 2012 (ADF&G 2017). Although the causes for this decline are not well documented, suitable freshwater habitat will be necessary for the conservation of this species into the future. We therefore recommend that ADEC temperature exceedances at the Lower Kelsall River and Jones Point sites especially be considered when making habitat management decisions in these waterbodies.

Furthermore, regular exceedances of ADEC standards at the Chilkat Weir during 2021 and 2022 should be examined further. The weir constricts passage to direct fish through a sonar counter when it is open, and blocks fish passage when it is closed at night. TWC is currently discussing investigation and mitigation strategies at the weir with ADF&G biologists.

With the help of our partners, TWC will continue to collect temperature and characteristic data at all of our sites, as well as establish multiple new sites within the Chilkat and Chilkoot watersheds. In 2023, we plan to also collect thermal imagery of the Chilkat watershed to identify thermal refugia, as well as areas of concern, with a high degree of resolution. In collaboration with The Southeast Alaska Freshwater Temperature Monitoring Network, we aim to generate a dataset that helps us understand thermal patterns and trends locally as well as regionally in order to help managers make effective decisions in response to climate change.

Literature Cited

- Alaska Department of Environmental Conservation (ADEC). 2020. 18 AAC 70, Water Quality Standards. Online: <https://dec.alaska.gov/water/water-quality/standards/>
- Alaska Department of Fish and Game (AGF&G). 2012. Subsistence Harvest and Data Reports. Online: <http://www.adfg.alaska.gov/sb/CSIS/index.cfm?ADFG=harvInfo.harvestCommSelComm>
- Alaska Department of Fish and Game (AGF&G). 2017. State of Alaska Special Status Species: Fish Stock of Concern. Online: <http://www.adfg.alaska.gov/index.cfm?adfg=specialstatus.akfishstocks>
- Alaska Department of Fish and Game (AGF&G). 2020. Anadromous Waters Catalog: Interactive Mapper. Online: <https://www.adfg.alaska.gov/sf/SARR/AWC/index.cfm?ADFG=main.interactive>
- Bellmore R., M. Winfree. Southeast Alaska Freshwater Monitoring Network Implementation Plan. 2019. Online: <https://www.alaskawatershedcoalition.org/southeast-alaska-stream-temperature-monitoring-network/>.
- Isaak D.J. and B.E. Rieman. 2013. Stream isotherm shifts from climate change and implications for distributions of ectothermic organisms. *Global Change Biol.* 19: 742–751.
- Kovach, R. P., J.E. Joyce, J.D. Echave, M.S. Lindberg, and D.A. Tallmon. (2013) Earlier Migration Timing, Decreasing Phenotypic Variation, and Biocomplexity in Multiple Salmonid Species, ed S M Carlson, *PLoS ONE*, 8 1–10 Online: <http://dx.plos.org/10.1371/journal.pone.0053807>.
- Mauger, S. 2013. Stream Temperature Monitoring for Cook Inlet Salmon Streams 2008-2012. Final report for the Alaska Department of Environmental Conservation and U.S. Fish and Wildlife Service. Online: <https://inletkeeper.org/wp-content/uploads/2017/03/Cook-Inlet-Stream-Temp-Network-Synthesis-Report.pdf>
- Mauger, S., R. Shaftel, E.J. Trammell, M. Geist, and D. Bogan. 2014. Stream temperature data collection standards and protocol for Alaska: minimum standards to generate data useful for regional-scale analyses. Cook Inletkeeper, Homer, AK and Alaska Natural Heritage Program, UAA, Anchorage, AK. 53 pp.
- Mauger, S., R. Shaftel, J.C. Leppi, and D.J. Rinella. 2017. Summer temperature regimes in southcentral Alaska streams: watershed drivers of variation and potential implications for Pacific salmon. *Can. J. Fish. Aquat. Sci.*, 74 702–15 Online: <http://www.nrcresearchpress.com/doi/10.1139/cjfas-2016-0076>.
- Schoen, E.R., M.S. Wipfli, E.J. Trammell, D.J. Rinella, A.J. Floyd, J. Grunblatt, M.D. McCarthy, B.E. Meyer, J. M. Morton, J.E. Powell, A. Prakash, M.N. Reimer, S.L. Stuefer, H. Toniolo, B.M. Wells, and F.D. Witmer. 2017. Future of pacific salmon in the face of environmental change: lessons from one of the World's remaining productive salmon regions. *Fisheries*, 42 (10) (2017). pp. 538-553
- Shanley, C. S., S. Pyare, M.I. Goldstein, P.B. Alaback, D.M. Albert, C.M. Beier, T.J. Brinkman, R.T. Edwards, E. Hood, A. MacKinnon, M.V. McPhee, T.M. Patterson, L.H. Suring, D.A. Tallmon, and M.S. Wipfli. 2015. Climatic change implications in the northern coastal temperate rainforest of North America. *Climatic Change*. 130: 155. Online: <https://doi.org/10.1007/s10584-015-1355-9>.

Steel, E. A., A. Tillotson, D.A. Larsen, A.H. Fullerton, K.P. Denton, and B.R. Beckman. 2012. Beyond the mean: The role of variability in predicting ecological effects of stream temperature on salmon, *Ecosphere*, 3 1–11 Online: <http://doi.wiley.com/10.1890/ES12-00255.1>.

Taylor, S. G. 2008. Climate warming causes phenological shift in Pink Salmon, *Oncorhynchus gorbuscha*, behavior at Auke Creek, Alaska: Climate warming and pink salmon behavior, *Glob. Change Biol.*, 14 229–35 Online: <http://doi.wiley.com/10.1111/j.1365-2486.2007.01494.x>.

Webb, B. W., D.M. Hannah, R.D. Moore, L.E. Brown, and F. Nobilis. 2008. Recent advances in stream and river temperature research, *Hydrol. Process.*, 22 902–18 Online: <http://doi.wiley.com/10.1002/hyp.6994>.



Chilkoot Lake. Photo by Derek Poinsette.